arXiv:1604.02712v2 [math.CO] 16 Aug 2016

On the number of mutually disjoint pairs of S-permutation matrices

Krasimir Yordzhev

Faculty of Mathematics and Natural Sciences
South-West University, Blagoevgrad, Bulgaria
E-mail: yordzhev@swu.bg

Abstract

This work examines the concept of S-permutation matrices, namely $n^2 \times n^2$ permutation matrices containing a single 1 in each canonical $n \times n$ subsquare (block). The article suggests a formula for counting mutually disjoint pairs of $n^2 \times n^2$ S-permutation matrices in the general case by restricting this task to the problem of finding some numerical characteristics of the elements of specially defined for this purpose factor-set of the set of $n \times n$ binary matrices. The paper describe an algorithm that solves the main problem. To do that, every $n \times n$ binary matrix is represented uniquely as a n-tuple of integers.

Keyword: binary matrix; S-permutation matrix; disjoint matrices; Sudoku; factor-set; n-tuple of integers

2010 Mathematics Subject Classification: 05B20

1 Introduction and notation

Let n be a positive integer. By $[n]$ we denote the set $[n] = \{1, 2, \ldots, n\}$.

A *binary* (or *boolean*, or (0,1)-*matrix*) is a matrix all of whose elements belong to the set $\mathfrak{B} = \{0, 1\}$. In this paper we will consider only square binary matrices. With $\mathfrak{B}_n$ we will denote the set of all $n \times n$ binary matrices. With $\mathfrak{B}_{n,k}$ we will denote the set of all $n \times n$ binary matrices containing exactly k elements equal to 1.

Two $n \times n$ binary matrices $A = (a_{ij}) \in \mathfrak{B}_n$ and $B = (b_{ij}) \in \mathfrak{B}_n$ will be called *disjoint* if there are not integers $i, j \in [n]$ such that $a_{ij} = b_{ij} = 1$, i.e. if $a_{ij} = 1$ then $b_{ij} = 0$ and if $b_{ij} = 1$ then $a_{ij} = 0$.

Let n be a positive integer and let $A \in \mathfrak{B}_{n^2}$ be a $n^2 \times n^2$ binary matrix. With the help of $n-1$ horizontal lines and $n-1$ vertical lines A has been separated into n^2 of number non-intersecting $n \times n$ square sub-matrices A_{kl}, $1 \leq k, l \leq n$, e.i.

$$A = \begin{bmatrix} A_{11} & A_{12} & \cdots & A_{1n} \\ A_{21} & A_{22} & \cdots & A_{2n} \\ \vdots & \vdots & \ddots & \vdots \\ A_{n1} & A_{n2} & \cdots & A_{nn} \end{bmatrix}. \tag{1}$$

The sub-matrices A_{kl}, $1 \leq k, l \leq n$ will be called *blocks*.

A matrix $A \in \mathfrak{B}_{n^2}$ is called an *S-permutation* if in each row, in each column, and in each block of A there is exactly one 1. Let the set of all $n^2 \times n^2$ S-permutation matrices be denoted by Σ_{n^2}.

The concept of S-permutation matrix was introduced by Geir Dahl [2] in relation to the popular Sudoku puzzle. Sudoku is a very popular game. On the other hand, it is well known that Sudoku matrices are special cases of Latin squares in the class of gerechte designs [1].

Obviously a square $n^2 \times n^2$ matrix M with elements of $[n^2] = \{1, 2, \ldots, n^2\}$ is a Sudoku matrix if and only if there are matrices $A_1, A_2, \ldots, A_{n^2} \in \Sigma_{n^2}$, each two of them are disjoint and such that P can be given in the following way:

$$M = 1 \cdot A_1 + 2 \cdot A_2 + \cdots + n^2 \cdot A_{n^2} \tag{2}$$

Some algorithms for obtaining random Sudoku matrices and their valuation are described in detail in [5] and [4].

In [4] Roberto Fontana offers an algorithm which randomly gets a family of $n^2 \times n^2$ mutually disjoint S-permutation matrices, where $n = 2, 3$. In $n = 3$ he ran the algorithm 1000 times and found 105 different families of nine mutually disjoint S-permutation matrices. Then using (2) he obtained $9! \cdot 105 = 38\ 102\ 400$ Sudoku matrices.

But it is known [3] that the total number of 9×9 Sudoku matrices is

$$9! \cdot 72^2 \cdot 2^7 \cdot 27\ 704\ 267\ 971 = 6\ 670\ 903\ 752\ 021\ 072\ 936\ 960$$

Thus, in relation with Fontana's algorithm, it looks useful to calculate the probability of two randomly generated S-permutation matrices to be disjoint. So the question of enumerating all disjoint pairs of S-permutation matrices naturally arises. This work is devoted to this task.

As we have shown in [6], with hand calculations of the so assigned task with small values of n ($n = 2, 3$), it is convenient to use the apparatus of graph theory. Unfortunately, when $n \geq 4$ this approach is inefficient. In this article, we will use only the operations of matrix analysis, which are not difficult to process with computers.

2 A representation of S-permutation matrices

Let n be a positive integer. If $z_1\ z_2\ \ldots\ z_n$ is a permutation of the elements of the set $[n] = \{1, 2, \ldots, n\}$ and let us shortly denote σ this permutation. Then in this case we will denote by $\sigma(i)$ the i-th element of this permutation, i.e. $\sigma(i) = z_i$, $i = 1, 2, \ldots, n$.

Definition 1 *Let Π_n denotes the set of all $n \times n$ matrices, constructed such that $\pi \in \Pi_n$ if and only if the following three conditions are true:*

i) *the elements of π are ordered pairs of numbers $\langle i, j \rangle$, where $1 \leq i, j \leq n$;*

ii) *if*

$$[\langle a_1, b_1 \rangle \quad \langle a_2, b_2 \rangle \quad \cdots \quad \langle a_n, b_n \rangle]$$

is the i-th row of π for any $i \in [n] = \{1, 2, \ldots, n\}$, then $a_1\ a_2\ \ldots\ a_n$ in this order is a permutation of the elements of the set $[n]$;

iii) *if*

$$\begin{bmatrix} \langle a_1, b_1\rangle \\ \langle a_2, b_2\rangle \\ \vdots \\ \langle a_n, b_n\rangle \end{bmatrix}$$

is the j-th column of π for any $j \in [n]$, then $b_1, b_2, \ldots, b_n$ in this order is a permutation of the elements of the set $[n]$.

From Definition 1, it follows that we can represent each row and each column of a matrix $M \in \Pi_n$ with the help of a permutation of elements of the set $[n]$.

Conversely for every $(2n)$-tuple

$$\langle\langle \rho_1, \rho_2, \ldots, \rho_n\rangle, \langle \sigma_1, \sigma_2, \ldots, \sigma_n\rangle\rangle,$$

where

$$\rho_i = \rho_i(1)\ \rho_i(2)\ \ldots\ \rho_i(n), \quad 1 \le i \le n$$
$$\sigma_j = \sigma_j(1)\ \sigma_j(2)\ \ldots\ \sigma_j(n), \quad 1 \le j \le n$$

are $2n$ permutations of elements of $[n]$ (not necessarily different), then the matrix

$$\pi = \begin{bmatrix} \langle \rho_1(1), \sigma_1(1)\rangle & \langle \rho_1(2), \sigma_2(1)\rangle & \cdots & \langle \rho_1(n), \sigma_n(1)\rangle \\ \langle \rho_2(1), \sigma_1(2)\rangle & \langle \rho_2(2), \sigma_2(2)\rangle & \cdots & \langle \rho_2(n), \sigma_n(2)\rangle \\ \vdots & \vdots & \ddots & \vdots \\ \langle \rho_n(1), \sigma_1(n)\rangle & \langle \rho_n(2), \sigma_2(n)\rangle & \cdots & \langle \rho_n(n), \sigma_n(n)\rangle \end{bmatrix}$$

is matrix of Π_n. Hence

$$|\Pi_n| = (n!)^{2n} \tag{3}$$

Definition 2 *We say that matrices $\pi' = \left[{p'}_{ij}\right]_{n\times n} \in \Pi_n$ and $\pi'' = \left[{p''}_{ij}\right]_{n\times n} \in \Pi_n$ are* disjoint*, if ${p'}_{ij} \neq {p''}_{ij}$ for every $i, j \in [n]$.*

Definition 3 *Let $\pi', \pi'' \in \Pi_n$, $\pi' = \left[{p'}_{ij}\right]_{n\times n}$, $\pi'' = \left[{p''}_{ij}\right]_{n\times n}$ and let the integers $i, j \in [n]$ are such that ${p'}_{ij} = {p''}_{ij}$. In this case we will say that ${p'}_{ij}$ and ${p''}_{ij}$ are* component-wise equal elements*.*

Obviously two Π_n-matrices are disjoint if and only if they do not have component-wise equal elements.

Example 1 We consider the following Π_3-matrices:

$$\pi' = \left[p'_{ij}\right] = \begin{bmatrix} \langle 3,1\rangle & \langle 2,1\rangle & \langle 1,2\rangle \\ \langle 2,3\rangle & \langle 3,2\rangle & \langle 1,1\rangle \\ \langle 3,2\rangle & \langle 1,3\rangle & \langle 2,3\rangle \end{bmatrix}$$

$$\pi'' = \left[p''_{ij}\right] = \begin{bmatrix} \langle 3,2\rangle & \langle 1,3\rangle & \langle 2,1\rangle \\ \langle 3,3\rangle & \langle 1,1\rangle & \langle 2,2\rangle \\ \langle 2,1\rangle & \langle 1,2\rangle & \langle 3,3\rangle \end{bmatrix}$$

$$\pi''' = \left[p'''_{ij}\right] = \begin{bmatrix} \langle 3,1\rangle & \langle 1,3\rangle & \langle 2,2\rangle \\ \langle 2,2\rangle & \langle 3,1\rangle & \langle 1,1\rangle \\ \langle 2,3\rangle & \langle 1,2\rangle & \langle 3,3\rangle \end{bmatrix}$$

Matrices π' and π'' are disjoint, because they do not have component-wise equal elements.

Matrices π' and π''' are not disjoint, because they have two component-wise equal elements: $p'_{11} = p'''_{11} = \langle 3, 1\rangle$ and $p'_{23} = p'''_{23} = \langle 1, 1\rangle$.

Matrices π'' and π''' are not disjoint, because they have three component-wise equal elements: $p''_{12} = p'''_{12} = \langle 1, 3\rangle$, $p''_{32} = p'''_{32} = \langle 1, 2\rangle$, and $p'_{33} = p'''_{33} = \langle 3, 3\rangle$.

The relationship between S-permutation matrices and the matrices from the set Π_n are given by the following theorem:

Theorem 1 *Let n be an integer, $n \geq 2$. Then there is one to one correspondence between the sets Σ_{n^2} and Π_n.*

Proof. Let $A \in \Sigma_{n^2}$. Then A is constructed with the help of formula (1) and for every $i, j \in [n]$ in the block A_{ij} there is only one 1 and let this 1 has coordinates (a_i, b_j). For every $i, j \in [n]$ we obtain ordered pairs of numbers $\langle a_i, b_j\rangle$ corresponding to these coordinates. As in every row and every column of A there is only one 1, then the matrix $[\alpha_{ij}]_{n\times n}$, where $\alpha_{ij} = \langle a_i, b_j\rangle$, $1 \leq i, j \leq n$, which is obtained by the ordered pairs of numbers is matrix of Π_n, i.e. matrix for which the conditions i), ii) and iii) are true.

Conversely, let $[\alpha_{ij}]_{n\times n} \in \Pi_n$, where $\alpha_{ij} = \langle a_i, b_j\rangle$, $i, j \in [n]$, $a_i, b_j \in [n]$. Then for every $i, j \in [n]$ we construct a binary $n \times n$ matrices A_{ij} with only one 1 with coordinates (a_i, b_j). Then we obtain the matrix of type (1). According to the properties i), ii) and iii), it is obvious that the obtained matrix is S-permutation matrix. □

Corollary 1 *The number of all pairs of disjoint matrices from Σ_{n^2} is equal to the number of all pairs of disjoint matrices from Π_n.*

Proof. It is easy to see that with respect of the described in Theorem 1 one to one correspondence, every pair of disjoint matrices of Σ_{n^2} will correspond to a pair of disjoint matrices of Π_n and conversely every pair of disjoint matrices of Π_n will correspond to a pair of disjoint matrices of Σ_{n^2}. □

Corollary 2 [2] *The number of all $n^2 \times n^2$ S-permutation matrices is equal to*

$$|\Sigma_{n^2}| = (n!)^{2n} \tag{4}$$

Proof. It follows immediately from Theorem 1 and formula (3). □

3 A formula for counting all disjoint pairs of $n^2 \times n^2$ S-permutation matrices

Let $A = [a_{ij}]_{n\times n} \in \mathfrak{B}_n$. We define the following numerical characteristics of the binary matrix A:

$r_k(A)$ – the number of rows in A having exactly k units, $k = 0, 1, 2, \ldots, n$;

$c_k(A)$ – the number of columns in A having exactly k units, $k = 0, 1, 2, \ldots, n$;

$\psi_k(A) = r_k(A) + c_k(A)$, $k = 0, 1, 2, \ldots, n$;

$\varepsilon(A)$ – the number of units in A.

Let $A, B \in \mathfrak{B}_n$. We will say that $A \sim B$, if B is obtained from A after dislocation of some of the rows of A. Obviously, the relation defined like that is an equivalence relation. The factor-set $\mathfrak{B}_{n/\sim}$, i.e. the set of equivalence classes on the above defined relation we denote with $\overline{\mathfrak{B}}_n$. If $A \in \mathfrak{B}_n$, then with $\overline{A}$ we will denote the set $\overline{A} = \{B \in \mathfrak{B}_n \mid B \sim A\}$. Thus $|\overline{A}| = |\{B \in \mathfrak{B}_n \mid B \sim A\}|$ is the cardinality of the set $\overline{A}$. By definition $\overline{\mathfrak{B}}_{n,k} = \mathfrak{B}_{n,k/\sim}$

Obviously if $A, B \in \mathfrak{B}_n$ and $A \sim B$, then $r_k(A) = r_k(B)$, $c_k(A) = c_k(B)$, $\psi_k(A) = \psi_k(B)$, $\varepsilon_k(A) = \varepsilon_k(B)$, $k = 0, 1, 2, \ldots, n$. So in a natural way we can define the functions r_k, c_k, ψ_k and ε in the factor-set $\overline{\mathfrak{B}}_n = \mathfrak{B}_{n/\sim}$ as $r_k(\overline{A})$, $c_k(\overline{A})$, $\psi_k(\overline{A})$ and $\varepsilon(\overline{A})$ will mean respectively $r_k(A)$, $c_k(A)$, $\psi_k(A)$ $\varepsilon(A)$, where A is an arbitrary representative of the set $\overline{A} = \{B \in \mathfrak{B}_n \mid B \sim A\}$.

Lemma 2 *Let $\pi \in \Pi_n$. Then the number $q(n,k)$ of all matrices $\pi' \in \Pi_n$ (including π), having at least k, $k = 0, 1, \ldots, n^2$ component-wise equal elements to the matrix π is equal to*

$$q(n,k) = \sum_{\overline{A} \in \overline{\mathfrak{B}}_{n,k}} |\overline{A}| \prod_{i=0}^{n-2} [(n-i)!]^{\psi_i(\overline{A})} \tag{5}$$

Proof. Let $\pi = [p_{ij}]_{n\times n}, \pi' = \left[p'_{ij}\right]_{n\times n} \in \Pi_n$ and let π and π' have exactly k component-wise equal elements. Then we uniquely obtain the binary $n \times n$ matrix $A = [a_{ij}]_{n\times n}$, such that $a_{ij} = 1$ if and only if $p_{ij} = p'_{ij}$, $i, j \in [n]$.

Inversely, let $A = [a_{ij}]_{n\times n} \in \mathfrak{B}_n$ and let $\pi = [p_{ij}]_{n\times n}$ be an arbitrary matrix from Π_n.We search for the number $h(\pi, A)$ of all matrices $\pi' = [p'_{ij}]_{n\times n} \in \Pi_n$, such that $p'_{ij} = p_{ij}$, if $a_{ij} = 1$. (It is assumed that there exist $s, t \in [n]$ such that $a_{st} = 0$ and $p'_{st} = p_{st}$.)

Let us denote with γ_s the number of 1 in s-th row of A and let the s-th row of π correspond to the permutation ρ_s of the elements of $[n]$, $s = 1, 2, \ldots, n$. Then there exist $(n-\gamma_s)!$ permutations ρ' of the elements of $[n]$, such that if $a_{st} = 1$, then $\rho_s(t) = \rho'(t)$, $t \in [n]$. Likewise we also prove the respective statement for the columns of π. Therefore

$$h(\pi, A) = \prod_{i=0}^{n} [(n-i)!]^{r_i(A)} \prod_{i=0}^{n} [(n-i)!]^{c_i(A)} = \prod_{i=0}^{n} [(n-i)!]^{\psi_i(A)}.$$

From everything said so far it follows that for each $\pi \in \Pi_n$ there exist

$$q(n,k) = \sum_{A \in \mathfrak{B}_{n,k}} \prod_{i=0}^{n} [(n-i)!]^{\psi_i(A)} = \sum_{\overline{A} \in \overline{\mathfrak{B}}_{n,k}} |\overline{A}| \prod_{i=0}^{n} [(n-i)!]^{\psi_i(\overline{A})}$$

matrices from Π_n, which have at least k elements that are component-wise equal to the respective elements of π.

And since $(n-n)! = 0! = 1$ and $[n-(n-1)]! = 1! = 1$, then we finally obtain formula (5). □

Lemma 3 *For every integer $n \geq 2$*

$$q(n,0) = q(n,1) = (n!)^{2n} = |\Pi_n| = |\Sigma_{n^2}|.$$

Proof. Let $k = 0$. Then $\mathfrak{B}_{n,0}$ contains only the matrix, all elements of which are equal to 0. So $|\mathfrak{B}_{n,0}| = 1$ and if $A \in \mathfrak{B}_{n,0}$ then $|\overline{A}| = 1$, $\psi_0(A) = 2n$ and $\psi_i(A) = 0$ when $i \geq 1$. Therefore $q(n,0) = \sum_{\overline{A} \in \overline{\mathfrak{B}}_{n,0}} |\overline{A}| \prod_{i=0}^{n-2} [(n-i)!]^{\psi_i(\overline{A})} = 1 \cdot [(n-0)!]^{2n} \prod_{i=1}^{n-2} [(n-i)!]^0 = (n!)^{2n}$.

When $k = 1$, there are n^2 matrices $A \in \mathfrak{B}_{n,1}$. It is easy to see that $|\overline{\mathfrak{B}}| = n$ and for every $\overline{A} \in \overline{\mathfrak{B}}_{n,1}$, $|\overline{A}| = n$, $\psi_0(\overline{A}) = 2(n-1)$, $\psi_1(\overline{A}) = 2$ and $\psi_i(\overline{A}) = 0$ for $i > 1$. Therefore $q(n,1) = n^2[(n-0)!]^{2n-2}[(n-1)!]^2 = (n!)^{2n-2}(n!)^2 = (n!)^{2n}$. □

Theorem 4 *Let $A \in \Sigma_{n^2}$. Then the number ξ_n of all matrices $B \in \Sigma_{n^2}$ which are disjoint with A does not depend on A and is equal to*

$$\xi_n = \sum_{\overline{A} \in \overline{\mathfrak{B}}_n,\ \varepsilon(\overline{A}) \geq 2} (-1)^{\varepsilon(\overline{A})} |\overline{A}| \prod_{i=0}^{n-2} [(n-i)!]^{\psi_i(\overline{A})} \tag{6}$$

Proof. Let $n \geq 2$ be an integer. Then applying Theorem 1, Lemma 2, Lemma 3 and the principle of inclusion and exclusion we obtain that the number ξ_n of all matrices $B \in \Sigma_{n^2}$ which are disjoint with A is equal to

$$\begin{aligned} \xi_n &= |\Pi_n| + \sum_{k=1}^{n^2} (-1)^k q(n,k) \\ &= (n!)^{2n} - (n!)^{2n} + \sum_{k=2}^{n^2} (-1)^k q(n,k) \\ &= \sum_{k=2}^{n^2} (-1)^k q(n,k), \end{aligned}$$

where the function $q(n,k)$ is calculated with the help of formula (5). Thus we obtain the proof to formula (6). □

Corollary 3 *The cardinality η_n of the set of all disjoint non-ordered pairs of $n^2 \times n^2$ S-permutation matrices is equal to*

$$\eta_n = \frac{(n!)^{2n}}{2} \xi_n \tag{7}$$

where ξ_n is described using formula 6.

Proof. It follows directly from formula (4) and having in mind that the "disjoint" relation is symmetric and antireflexive. □

Corollary 4 *The probability p_n of two randomly generated $n^2 \times n^2$ S-permutation matrices to be disjoint is equal to*

$$p_n = \frac{\xi_n}{(n!)^{2n} - 1}, \tag{8}$$

where ξ_n is described using formula (6).

Proof. Applying Corollary 3 and formula (4), we obtain:

$$p_n = \frac{\eta_n}{\binom{|\Sigma_{n^2}|}{2}} = \frac{\frac{(n!)^{2n}}{2}\xi_n}{\frac{(n!)^{2n}\left((n!)^{2n}-1\right)}{2}} = \frac{\xi_n}{(n!)^{2n}-1}.$$

□

4 An algorithm for counting

There is one to one correspondence between the representation of the integers in decimal and in binary notations. So a square binary $n \times n$ matrix can be represented using ordered n-tuple of nonnegative integers, which belong to the closed interval $[0,\ 2^n-1]$. Let the integer $a \in [0,\ 2^n-1]$. Then a is represented uniquely in the form:

$$a = \sum_{u=0}^{n-1} b_u(a)2^u,$$

where $b_u(a) \in \mathfrak{B} = \{0,1\},\ u = 0,1,\ldots,n-1$. We assume that we have implemented an algorithm for calculating the functions $b_u(a)$ for every $u = 0,1,\ldots,n-1$ and for every $a \in [0,\ 2^n-1]$. For example, in the programming languages C ++ and Java, $b_u(a)$ can be calculated using the expression

```
bu = (a & (1<<u))==0 ? 0 : 1
```

Let $A \in \mathfrak{B}_n$. With $\rho(A)$ we will denote the ordered n-tuple

$$\rho(A) = \langle x_1, x_2, \ldots, x_n\rangle,$$

where $0 \le x_i \le 2^n-1,\ i = 1,2,\ldots n$ and x_i is the integer written in binary notation with the help of the i-th row of A.

We consider the set:

$$\begin{aligned}\mathfrak{R}_n &= \{\langle x_1, x_2, \ldots, x_n\rangle \mid 0 \le x_i \le 2^n-1,\ i = 1,2,\ldots n\}\\ &= \{\rho(A) \mid A \in \mathfrak{B}_n\}\end{aligned}$$

Thus we define the mapping $\rho : \mathfrak{B}_n \to \mathfrak{R}_n$, which is bijective and therefore $\mathfrak{B}_n \cong \mathfrak{R}_n$.

If $A \in \mathfrak{B}_n$ and $\rho(A) = \alpha \in \mathfrak{R}_n$, then by analogy we define the numerical characteristics of the element $\alpha \in \mathfrak{R}_n$: $r_k(\alpha) = r_k(A)$, $c_k(\alpha) = c_k(A)$, $\psi_k(\alpha) = r_k(\alpha)+c_k(\alpha) = \psi_k(A)$, $k = 0,1,2,\ldots,n$ and $\varepsilon(\alpha) = \varepsilon(A)$. We assume $|\alpha| = |\overline{A}|$, where $\overline{A} = \{B \in \mathfrak{B}_n \mid B \sim A\}$.

Let $\alpha = \langle x_1, x_2, \ldots, x_n\rangle \in \mathfrak{R}_n$ and let s be the number of different elements in $\alpha = \langle x_1, x_2, \ldots, x_n\rangle$. Then the set $X = \{x_1, x_2, \ldots, x_n\}$ can be divide into parts

$$X = X_1 \cup X_2 \cup \cdots \cup X_s$$

such that for every $k \in [s]$ and every $i,j \in [n]$, $i \ne j$ the condition $x_i, x_j \in X_k$ is satisfied if and only if $x_i = x_j$. We assume

$$z_i = |X_i|, \quad i = 1,2,\ldots s.$$

It is easily seen that

$$|\alpha| = \frac{n!}{\prod_{i=1}^{s} z_i!}.$$

Let

$$\overline{\mathfrak{R}}_n = \{\langle x_1, x_2, \ldots, x_n\rangle \mid 0 \le x_1 \le x_2 \le \cdots \le x_n \le 2^n - 1\} \subset \mathfrak{R}_n.$$

It is easily seen that $\overline{\mathfrak{B}}_n \cong \overline{\mathfrak{R}}_n$, which gives the basis to construct the following algorithm for calculating ξ_n:

Algorithm 5 *Calculation of* ξ_n.
begin
$\xi_n := 0$;
For every $\alpha = \langle x_1 x_2, \ldots, x_n\rangle \in \overline{\mathfrak{R}}_n$ *do*
{
$s := 1$;
$\varepsilon(\alpha) := 0$;
For $i = 1, 2, \ldots, n$ *do*
{
$z_s := z_s + 1$;
$t = 0$;
For $u = 0, 1, \ldots, n-1$ *do*
{
$t := t + b_u(x_i)$;
}
$r_t(\alpha) := r_t(\alpha) + 1$;
$\varepsilon(\alpha) := \varepsilon(\alpha) + t$;
If $i < n$ *and* $x_i < x_{i+1}$ *then* $s := s + 1$;
}
If $\varepsilon(\alpha) = 0$ *or* $\varepsilon(\alpha) = 1$ *then go to next* α;
For $u = 0, 1, \ldots, n-1$ *do*
{
$t := 0$;
For $i = 1, 2, \ldots, n$ *do*
{
$t = t + b_u(x_i)$;
}
$c_t(\alpha) := c_t(\alpha) + 1$;
}
For $k = 0, 1, \ldots, n$ *do*
{
$\psi_k(\alpha) := r_k(\alpha) + c_k(\alpha)$;
}
$|\alpha| := \dfrac{n!}{\prod_{i=1}^{s} z_i!}$;
$T(\alpha) := (-1)^{\varepsilon(\alpha)} |\alpha| \prod_{i=0}^{n-2} [(n-i)!]^{\psi_i(\alpha)}$;

$\xi_n := \xi_n + T(\alpha)$;
}
end.

5 Conclusion

On the basis of algorithm 5 with programming language Java, we made a computer program for calculating ξ_n, η_n and p_n and we received the following results:

$$\xi_2 = 7$$
$$\xi_3 = 17\ 972$$
$$\xi_4 = 41\ 685\ 061\ 617$$
$$\xi_5 = 232\ 152\ 032\ 603\ 580\ 176\ 504$$
$$\xi_6 = 7\ 236\ 273\ 578\ 711\ 450\ 275\ 537\ 707\ 547\ 657\ 855$$

$$\eta_2 = 56$$
$$\eta_3 = 419\ 250\ 816$$
$$\eta_4 = 2\ 294\ 248\ 126\ 968\ 596\ 791\ 296$$
$$\eta_5 = 71\ 871\ 209\ 790\ 288\ 983\ 974\ 921\ 874\ 964\ 480\ 000\ 000\ 000$$
$$\eta_6 = 7\ 022\ 228\ 210\ 556\ 132\ 949\ 916\ 635\ 069\ 726\ 824\ 032\ 981\ 704\ 989\ 720\ 182\ 784 \cdot 10^{13}$$

$$p_2 = 0.4666666666666667$$
$$p_3 = 0.38521058836137606$$
$$p_4 = 0.3786958223051558$$
$$p_5 = 0.37493849344703684$$
$$p_6 = 0.3728421644517476$$

For n = 2 and n = 3, the results that we get here coincide with the calculations made by hand in [6], where we used a graph theory approach.

References

[1] R.A. Bailey, P.J. Cameron, and R. Connelly. Sudoku, gerechte designs, resolutions, affine space, spreads, reguli, and hamming codes. *Amer. Math. Monthly*, (115):383–404, 2008.

[2] Geir Dahl. Permutation matrices related to sudoku. *Linear Algebra and its Applications*, 430(8–9):2457–2463, 2009.

[3] Bertram Felgenhauer and Frazer Jarvis. Enumerating possible sudoku grids, 2005. http://www.afjarvis.staff.shef.ac.uk/sudoku/sudoku.pdf.

[4] Roberto Fontana. Fractions of permutations. an application to sudoku. *Journal of Statistical Planning and Inference*, 141(12):3697–3704, 2011.

[5] Krasimir Yordzhev. Random permutations, random sudoku matrices and randomized algorithms. *International J. of Math. Sci. & Engg. Appls.*, 6(VI):291 – 302, 2012.

[6] Krasimir Yordzhev. Calculation of the number of all pairs of disjoint s-permutation matrices. *Applied Mathematics and Computation*, 268:1 – 11, 2015.